

4-Phenyl-1,2,4-triazoline-3,5-dione in the ene reactions with cyclohexene, 1-hexene and 2,3-dimethyl-2-butene. the heat of reaction and the influence of temperature and pressure on the reaction rate

Kiselev V., Kornilov D., Kashaeva H., Potapova L., Konovalov A.
Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia

Abstract

The values of the enthalpy (53.3; 51.3; 20.0 kJ mol⁻¹), entropy (-106; -122; -144 J mol⁻¹K⁻¹), and volume of activation (-29.1; -31.0; -cm³ mol⁻¹), the reaction volume (-25.0; -26.6; -cm³ mol⁻¹) and reaction enthalpy (-155.9; -158.2; -150.2 kJ mol⁻¹) have been obtained for the first time for the ene reactions of 4-phenyl-1,2,4-triazoline-3,5-dione 1, with cyclohexene 4, 1-hexene 6, and with 2,3-dimethyl-2-butene 8, respectively. The ratio of the values of the activation volume to the reaction volume ($\Delta V^\ddagger_{\text{corr}}/\Delta V_r - n$) in the ene reactions under study, 1 + 4 → 5 and 1 + 6 → 7, appeared to be the same, namely 1.16. The large negative values of the entropy and the volume of activation of studied reactions 1 + 4 → 5 and 1 + 6 → 7 better correspond to the cyclic structure of the activated complex at the stage determining the reaction rate. The equilibrium constants of these ene reactions can be estimated as exceeding 10¹⁸ L mol⁻¹, and these reactions can be considered irreversible. Copyright © 2014 John Wiley & Sons, Ltd.

<http://dx.doi.org/10.1002/poc.3277>

Keywords

1-hexene, 2,3-dimethylbutene-2, 4-phenyl-1,2,4-triazolin-3,5-dione, activation volume, cyclohexene, ene-reaction, enthalpy reaction, high pressure effect, rate constant